

Factors Associated with Burnout Syndrome among Clinical Medical Students at Chonburi Hospital, Thailand

Supitcha Thamissarakul, M.D.¹, Wasana Hongkan, M.D.¹, Prakasit Wannapaschaiyong, M.D.²

¹Chonburi Hospital Medical Education Center, Chonburi Hospital, Mueang, Chonburi 20000, Thailand.

²Department of Pediatrics, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkoknoi, Bangkok 10700, Thailand.

Received 4 August 2023 • Accepted 28 August 2023 • Published online 26 December 2023

Abstract:

Objective: Burnout syndrome causes chronic fatigue, loss of motivation, despair, decreased work performance, and depression. Clinical medical students risk developing burnout syndrome due to long-term learning pressure, responsibilities for patient care and night shifts. This study aimed to examine the factors associated with burnout syndrome among clinical medical students at Chonburi Hospital.

Material and Methods: A cross-sectional study was conducted from July 2022 and January 2023. Data were collected using the Maslach Burnout Inventory General Survey among 106 medical students. Descriptive statistics, chi-square, ANOVA, and the independent t-test were used to investigate the association among variables.

Results: Severe burnout syndrome was not found among clinical medical students. High emotional exhaustion was the most common burnout symptom among these medical students (53.8%). While 27.3% had high depersonalization, no medical students had low personal accomplishment. The factor related to high emotional exhaustion was 4th-year medical students. At the same time, the factors associated with high depersonalization were fourth-year medical students, males, and low accumulated grade point averages (GPAX).

Conclusion: The prevalence of severe burnout was not found among clinical medical students at Chonburi Hospital. However, males, fourth-year medical students, and low GPAX were risk factors for burnout symptoms. Therefore, screening, monitoring, and providing help to these at-risk medical students to reduce their chances of severe burnout should be encouraged.

Keywords: burnout, clinical medical students, depersonalization, emotional exhaustion

Contact: Prakasit Wannapaschaiyong, M.D.

Division of Developmental and Behavioral Pediatrics, Department of Pediatrics,
Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkoknoi, Bangkok 10700, Thailand.
E-mail: pra_ka_sit@hotmail.com

J Health Sci Med Res

doi: 10.31584/jhsmr.20231020

www.jhsmr.org

© 2023 JHSMR. Hosted by Prince of Songkla University. All rights reserved.

This is an open access article under the CC BY-NC-ND license

(<http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy>).

Introduction

Burnout syndrome is a psychological alteration caused by a response to chronic work stress or interpersonal stress¹. Prevalent in healthcare providers, this condition can cause them to experience both physical and mental exhaustion, leading to chronic fatigue, headaches, loss of motivation, and hopelessness². Suppose that people with this condition do not receive proper treatment they can develop depression and insomnia, resulting in frequent absenteeism and decreased work productivity³. Generally, burnout syndrome is assessed based on three main symptoms: emotional exhaustion (EE), or feeling overwhelmed with emotional pressure and lack of energy to face work; depersonalization (DP), defined as having a negative attitude toward work, resulting in an indifferent response to work; and reduced personal accomplishment (PA), which refers to decreased self-efficacy and efficiency at work⁴.

According to previous studies, the prevalence of burnout syndrome was as high as 45–56% among medical students^{5,6}. When considering each problem, almost half of all medical students had a high level of EE, and one-third had a high level of DP⁷. Factors related to burnout syndrome among medical students include females⁸, low academic performance⁹, practice on the wards of major subjects¹⁰, lack of motivation, difficulty adapting to medical learning styles and a desire to drop out¹¹.

Clinical medical students change their learning style, from studying in a preclinical classroom to practicing in the wards. Working in the wards, writing patient reports and working night shifts have resulted in stress from learning adjustments, maintaining relationships with people on the job, adapting to sleep deprivation and a lack of free time. These accumulate into continuous stress and cause a deterioration in their quality of life; additionally, the problem can affect their mental health and cause psychiatric disorders. Therefore, there is interest in studying the

prevalence and risk factors of burnout syndrome in clinical medical students at Chonburi Hospital, so as to find ways to prevent this condition.

Material and Methods

Study design and population

This was a cross-sectional study among clinical medical students practicing at Chonburi Hospital, Thailand from July 2022 to April 2023. One hundred and six medical students participated in this study and completed the questionnaire. This study protocol was approved by the Chonburi Hospital Ethics Committee (code: 72/66/S/q).

Data collection

All clinical medical students practicing at Chonburi Hospital during the data collection period were included in this study. The objectives of the study were informed, and participants were provided assurances that all who participated would remain anonymous, in addition to all responses to the questionnaire would be kept confidential. This study used two questionnaires: a demographic information form and the Maslach Burnout Inventory-General Survey.

Research instruments

1. Demographic information form

This form recorded demographic characteristics including gender, cumulative grade point average (GPAX), and year level of medical students.

2. Maslach Burnout Inventory – General Survey (Thai version)¹²

This questionnaire is a standardized tool used to assess burnout syndrome. The self-reported questionnaire consisted of 22 items, rated on a 7-point Likert scale: 0 (never felt it) to 6 (feel it every day). The results are presented with respect to the high, moderate, and low severity of 3 core burnout symptoms: emotional exhaustion

(EE), depersonalization (DP), and personal accomplishment (PA). People with high EE, high DP, and low PA are considered to have severe burnout syndrome. This questionnaire had an overall Cronbach's alpha greater than 0.75.

Data analysis

Descriptive statistics were used to describe the percentage and mean \pm S.D. Fisher's exact test, the independent t-test, and ANOVA were used to analyze the association between demographic characteristics and symptoms of burnout. A p-value of <0.05 was considered statistically significant. All analysis were performed using IBM Statistical Package for the Social Sciences (SPSS) Statistics (version 25; IBM Corporation, Armonk, New York, USA).

Results

One hundred and six clinical medical students from Chonburi Hospital participated in this study (Table 1). More than half of the participants were male (53.8%). Participants were 22 to 24 years old, 33.0% in fourth-year, 34.9% in fifth-year, and 32.1% in sixth-year. More than half of the participants had a GPAX of ≥ 3.00 (53.8%). No participants with severe burnout syndrome were found.

Table 2 shows that clinical medical students had high levels of EE up to 53.8%. Meanwhile, nearly half of the clinical medical students (42.5%) had moderate DP symptoms. No clinical medical students with low levels of PA were found.

However, considering each year-level, it was found that fourth-year medical students had a significantly higher mean EE score than other years (p-value=0.015), and a higher proportion of fourth-year medical students had high EE levels (68.6%), comparing to other years. In the DP domain, the average score of fourth-year medical students was significantly higher than other years (p-value=0.042).

Additionally, higher levels of DP (40%) were found in fourth-year medical students than other years.

Taking into account gender, men were associated with higher DP scores and a more significant proportion of high DP symptoms than women (p-value=0.002 and 0.002, respectively) (Table 3). At the same time, lower GPAX was significantly associated with a higher proportion of high DP symptoms (p-value=0.048); as shown in Table 4.

Table 1 Baseline characteristics of clinical medical students (N=106)

Demographic characteristics	Descriptive results
Gender	
Male	57(53.8)
Female	49 (46.2)
Age (years) ^a	22.1 \pm 0.4
Year level of clinical medical students	
4 th year medical students	35 (33.0)
5 th year medical students	37 (34.9)
6 th year medical students	34 (32.1)
GPAX	
<3.00	49 (46.2)
≥ 3.00	57 (53.8)
Burnout syndrome	
Severe	0 (0.0)
Mild to moderate	106 (100)

GPAX=cumulative grade point average

Data presented as a number (percentage)

^aData presented as mean \pm standard deviation

Discussion

This study did not detect severe burnout among clinical medical students at Chonburi Hospital. These results are inconsistent with the study by Auchayasawat (2022), in which the prevalence of severe burnout syndrome among clinical medical students from Ubon Ratchathani University was as high as 31%¹⁰. At the same time, a meta-analysis of the global prevalence of medical students found that severe burnout was as high as 44.2%¹³. This difference may

be attributed to Chonburi Hospital's preventive measures against stress-related problems during medical student practice; such as arranging consultants to provide consulting services and organizing activities to encourage medical students to practice adaptive coping strategies to reduce the adverse effects of stress that can lead to severe burnout.

Considering all aspects of burnout syndromes, this study's participants had a high EE level of 53.8%, higher than other burnout symptoms. This result is consistent with

a study by Frajerma et al. (2019), which found that clinical medical students had emotional exhaustion as their most prevalent problem¹³. Increased stress from heavy workloads and responsibilities, the anticipation of medical staff, studies, and examinations that are relatively more difficult than the preclinical level may be the reasons why clinical medical students feel easily exhausted with work and service to patients¹⁴.

Table 2 Burnout symptoms of clinical medical students (N=106)

Burnout symptoms	Low level	Moderate level	High level	Average score ^a
EE	29 (27.3)	20 (18.9)	57 (53.8)	25.8±11.1
DP	32 (30.2)	45 (42.5)	29 (27.3)	9.8±4.9
PA	0 (0.0)	9 (8.5)	97 (91.5)	18.7±8.1

EE=emotional exhaustion, DP=depersonalization, PA=personal accomplishment

Data presented as a number (percentage), ^aData presented as mean±standard deviation

Table 3 Comparing the symptoms of burnout syndrome among 3 year-levels of clinical medical students

Symptoms of burnout syndrome	4 th year (n=35)	5 th year (n=37)	6 th year (n=34)	p-value
EE scores ^{a,b}	30.3±10.3	23.4±11.2	23.7±11.8	0.015*
Low level	5 (14.3)	15 (40.5)	9 (26.5)	0.121
Moderate level	6 (17.1)	7 (18.9)	7 (20.6)	
High level	24 (68.6)	15 (40.5)	18 (52.9)	
DP scores ^{a,b}	11.5±6.0	8.1±5.7	10.0±5.3	0.042*
Low level	7 (20)	17 (45.9)	8 (23.5)	0.038*
Moderate level	14 (40)	12 (32.4)	19 (55.9)	
High level	14 (40)	8 (21.6)	7 (20.6)	
PA scores ^{a,b}	19.4±7.6	18.9±8.7	17.7±7.7	0.701
Low level	0 (0.0)	0 (0.0)	0 (0.0)	0.758
Moderate level	3 (8.6)	4 (10.8)	2 (5.9)	
High level	32 (91.4)	33 (89.2)	32 (94.1)	

EE=emotional exhaustion, DP=depersonalization, PA=personal accomplishment

Data presented as number (percentage), ^aData presented as mean±standard deviation

The comparison of differences were analyzed by chi-square test and ^bANOVA

*Statistically significant at a p-value<0.05

Table 4 Comparison of the symptoms of burnout syndrome and their association with gender and GPAX

Symptoms of burnout syndrome	Male (n=57)	Female (n=49)	p-value	GPAX<3.00 (n=49)	GPAX≥3.00 (n=57)	p-value
EE scores ^{a,b}	27.5±12.52	27.2±10.38	0.907	28.3±12.2	26.6±10.9	0.439
Low level	14 (24.6)	9 (18.4)	0.374	11 (22.4)	12 (21.1)	0.655
Moderate level	11 (19.3)	15 (30.6)		10 (20.4)	16 (28.1)	
High level	32 (56.1)	25 (51.0)		28 (57.1)	29 (50.9)	
DP scores ^{a,b}	11.1±6.4	7.3±5.7	0.002*	10.6±6.6	8.3±6.0	0.064
Low level	13 (22.8)	26 (53.1)	0.002*	16 (32.7)	23 (40.4)	0.048*
Moderate level	21 (36.8)	15 (30.6)		13 (26.5)	23 (40.4)	
High level	23 (40.4)	8 (16.3)		20 (40.8)	11 (19.3)	
PA scores ^{a,b}	18.4±9.2	17.2±7.9	0.471	18.5±8.9	17.3±8.3	0.485
Low level	1 (1.8)	0 (0.0)	0.419	0 (0.0)	1 (1.8)	0.419
Moderate level	3 (5.3)	5 (10.2)		5 (10.2)	3 (5.3)	
High level	53 (93)	44 (89.8)		44 (89.8)	53 (93.0)	

EE=emotional exhaustion, DP=depersonalization, PA=personal accomplishment, GPAX=cumulative grade point average

Data presented as number (percentage), ^aData presented as mean±S.D.

The comparison of differences were analyzed by chi-square test and ^bindependent t-test

*Statistically significant at a p-value<0.05

This study found that fourth-year medical students had the highest proportion of high levels of EE and DP compared to other grades. The results of this study are similar to those of the Srikanth (2014)¹⁵ and Martini (2004)¹⁶ studies, which found that emotional exhaustion and depersonalization of residents in the first year were greater than that of residents in the second and third years. This result may be explained by the observation that those with more knowledge and work experience can better cope with stress and be more attentive.

In this study, men were associated with a high level of depersonalization, which is consistent with Korkate's study (2019)¹¹; as struggling men usually do not express or seek help¹⁷. For this reason, it can lead men to avoid facing pressure and neglecting others, which may explain this association with depersonalization.

Furthermore, a lower GPAX was associated with a high level of depersonalization, which is consistent with the studies by Srikanth (2014)¹⁵ and Apiwatanasiri (2007)⁹

these found that medical students with a lower grade point average were statistically significantly related to high levels of depersonalization. Medical students having insufficient knowledge can lead to insecurity in patient care and negative attitudes towards providing and responding to patients.

However, this study did not investigate many other factors related to burnout symptoms including current working environment, physical and mental health issues, financial problems, and attitudes toward studying medicine. Furthermore, this cross-sectional study design limits itself from concluding the cause and effect. Future longitudinal research should be conducted to determine the causes and investigate other burnout-related variables. Furthermore, the effectiveness of the implementation of the policy on stress prevention policy related to work in medical students at Chonburi Hospital should be studied, so as to benefit other medical schools in burnout syndrome prevention in medical students.

Conclusion

This study did not find severe burnout syndrome in clinical medical students at Chonburi Hospital. However, 4th-year medical students had higher emotional exhaustion and depersonalization. Male and a lower GPAX were related to depersonalization. Therefore, medical personnel should be encouraged to pay close attention to medical students with burnout risk factors. Additionally, policies should be formulated to prevent work-related stress to reduce negative consequences in the future.

Conflict of interest

All authors declare no conflict of interest.

References

1. Panagioti M, Geraghty K, Johnson J, Zhou A, Panagopoulou E, Chew-Graham C, et al. Association between physician burnout and patient safety, professionalism, and patient satisfaction: a systematic review and meta-analysis. *JAMA Intern Med* 2018; 178:1317–31.
2. Rotenstein LS, Torre M, Ramos MA, Rosales RC, Guille C, Sen S, et al. Prevalence of burnout among physicians: a systematic review. *JAMA* 2018;320:1131–50.
3. Wood BD, Killion JB. Burnout among healthcare professionals. *Radiol Manage* 2007;29:30–8.
4. Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol* 2001;52:397–422.
5. Dyrbye LN, West CP, Satele D, Boone S, Tan L, Sloan J, et al. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med* 2014;89:443–51.
6. Dyrbye LN, Thomas MR, Huntington JL, Lawson KL, Novotny PJ, Sloan JA, et al. Personal life events and medical student burnout: a multicenter study. *Acad Med* 2006;81:374–84.
7. Dyrbye L, Shanafelt T. A narrative review on burnout experienced by medical students and residents. *Med Educ* 2016;50:132–49.
8. Prins JT, Gazendam-Donofrio SM, Tubben BJ, van der Heijden FM, van de Wiel HB, Hoekstra-Weebers JE. Burnout in medical residents: a review. *Med Educ* 2007;41:788–800.
9. Apiwatanasiri C, Somakettrarin K, Suraprayoon K, Leurmpasert K, Wankaew N, Homchampa P. Stress and coping in medical students at clinical level, Khon Kaen University. *Srinagrind Med J* 2007;22:416–24.
10. Auchayasawat S, Parichart SP, Praken T, Ritdejch N, Yingsanongchat P, Usaneengamcharoen P. Prevalence and associated factors of burnout syndrome among clinical medical students. *J Psychiatr Assoc Thailand* 2022;67:101–12.
11. Korkate T, Krongkhwan R, Nirucha P, Winitra N. Prevalence and associated factors of burnout syndrome in preclinical medical students. *Thammasat Med J* 2019;19:127–38.
12. Summawart S. Burnout among the staff nurses in Ramathibodi Hospital [Master's Thesis]. Bangkok: Mahidol University; 1989.
13. Frajerma A, Morvan Y, Krebs MO, Gorwood P, Chaumette B. Burnout in medical students before residency: a systematic review and meta-analysis. *Eur Psychiatry* 2019;55:36–42.
14. Wannapaschaiyong P, Kallawicha K. Stress and stress coping strategies among foreign bachelor's medical interns in Bangkok Metropolitan Administration (BMA) hospitals. *J Health Res* 2023; 37:306–16.
15. Srikam S, Jiamjarasrangsri W, Lalitanantpong D. Job burnout and related factors among residents of King Chulalongkorn Memorial Hospital. *J Psychiatr Assoc Thailand* 2014;59:139–50.
16. Martini S, Arfken CL, Churchill A, Balon R. Burnout comparison among residents in different medical specialties. *Acad Psychiatry* 2004;28:240–2.
17. Houkes I, Winants Y, Twellaar M, Verdonk P. Development of burnout over time and the causal order of the three dimensions of burnout among male and female GPs. A three-wave panel study. *BMC Public Health* 2011;11:240.